

# Prevention and reversal of cardiac hypertrophy by soluble epoxide hydrolase inhibitors

Danyan Xu<sup>\*†</sup>, Ning Li<sup>\*</sup>, Yuxia He<sup>\*</sup>, Valeriy Timofeyev<sup>\*</sup>, Ling Lu<sup>\*</sup>, Hsing-Ju Tsai<sup>‡</sup>, In-Hae Kim<sup>‡</sup>, Dipika Tuteja<sup>\*</sup>, Robertino Karlo P. Mateo<sup>\*</sup>, Anil Singapuri<sup>\*</sup>, Benjamin B. Davis<sup>§</sup>, Reginald Low<sup>\*</sup>, Bruce D. Hammock<sup>\*¶</sup>, and Nipavan Chiamvimonvat<sup>\*¶\*\*</sup>

<sup>\*</sup>Division of Cardiovascular Medicine, <sup>§</sup>Center for Health and the Environment, and <sup>‡</sup>Department of Entomology and Cancer Research Center, University of California, Davis, CA 95616; <sup>¶</sup>Department of Veterans Affairs, Northern California Health Care System, Mather, CA 95655; and <sup>†</sup>Department of Cardiology, Internal Medicine, Xiangya Second Hospital, Central-South University, Changsha, Hunan Province 410007, China

Contributed by Bruce D. Hammock, October 17, 2006 (sent for review June 19, 2006)

Sustained cardiac hypertrophy represents one of the most common causes leading to cardiac failure. There is emerging evidence to implicate the involvement of NF- $\kappa$ B in the development of cardiac hypertrophy. However, several critical questions remain unanswered. We tested the use of soluble epoxide hydrolase (sEH) inhibitors as a means to enhance the biological activities of epoxyeicosatrienoic acids (EETs) to treat cardiac hypertrophy. sEH catalyzes the conversion of EETs to form the corresponding dihydroxyeicosatrienoic acids. Previous data have suggested that EETs may inhibit the activation of NF- $\kappa$ B-mediated gene transcription. We directly demonstrate the beneficial effects of several potent sEH inhibitors (sEHIs) in cardiac hypertrophy. Specifically, we show that sEHIs can prevent the development of cardiac hypertrophy using a murine model of pressure-induced cardiac hypertrophy. In addition, sEHIs reverse the preestablished cardiac hypertrophy caused by chronic pressure overload. We further demonstrate that these compounds potentially block the NF- $\kappa$ B activation in cardiac myocytes. Moreover, by using *in vivo* electrophysiologic recordings, our study shows a beneficial effect of the compounds in the prevention of cardiac arrhythmias that occur in association with cardiac hypertrophy. We conclude that the use of sEHIs to increase the level of the endogenous lipid epoxides such as EETs may represent a viable and completely unexplored avenue to reduce cardiac hypertrophy by blocking NF- $\kappa$ B activation.

epoxyeicosatrienoic acids | NF- $\kappa$ B

Cardiovascular disease represents an important and growing public health problem and is the cause of substantial morbidity and mortality. Cardiac hypertrophy is the heart's compensatory response to a variety of extrinsic and intrinsic stimuli including pressure or volume overload, mutations of sarcomeric proteins, or loss of contractile mass from previous myocardial infarction. Cardiac hypertrophy is believed to have a compensatory function by diminishing wall stress. Conversely, sustained cardiac hypertrophy represents one of the most common causes of cardiac failure (1, 2).

There is increasing evidence that prevention of cardiac hypertrophy may represent a new therapeutic target to improve patient survival (1, 3, 4). Inhibition of NF- $\kappa$ B has recently been investigated as one of the possible therapeutic approaches to treatment of cardiac hypertrophy. The results of the investigations may have important clinical implication (5–12). We surmised that the use of soluble inhibitors of NF- $\kappa$ B may present an attractive therapeutic means to treat cardiac hypertrophy. However, NF- $\kappa$ B can be both anti- and proapoptotic under certain conditions (13, 14). Therefore, the beneficial effects of NF- $\kappa$ B inhibitors in cardiac hypertrophy remain to be systematically investigated. Motivated by the uncertainties and the therapeutic potential of the inhibitors of this nuclear factor we tested the beneficial effects of several potent soluble epoxide hydrolase inhibitors (sEHIs) in cardiac hypertrophy. sEH catalyzes the conversion of epoxyeicosatrienoic acids (EETs) to form the corresponding dihydroxyeicosatrienoic acids (DHETs) (15, 16). EETs are products of cytochrome P450 epoxygenases that have vasodilatory properties similar to those of endothelium-

derived hyperpolarizing factor (17). In addition, EETs can inhibit the activation of NF- $\kappa$ B-mediated gene transcription (18, 19). Because sEH converts EETs to DHETs, inhibition of sEH has been intensely investigated as a potential approach for enhancing the antihypertensive and antiinflammatory actions of EETs (17, 20–22). However, the potential benefits of sEHIs and EETs for cardiac hypertrophy have never been investigated.

Here, we tested the effects of sEHIs on the development and reversal of cardiac hypertrophy using a murine model of aortic banding. We show that there is an almost complete resolution of cardiac hypertrophy by sEHIs independent of the antihypertensive effects. We further demonstrate that these compounds potentially block the NF- $\kappa$ B activation in cardiac myocytes. Moreover, our study shows a beneficial effect of sEHIs in the prevention of cardiac arrhythmias that occur in association with cardiac hypertrophy.

## Results

**sEHIs Prevent the Development of Cardiac Hypertrophy in a Murine Model with Thoracic Aortic Constriction (TAC).** We created a TAC in 8-week-old male C57BL/6J mice (Charles River Laboratories, Wilmington, MA), resulting in a chronic pressure overload-induced cardiac hypertrophy. TAC mice were randomized to receive either vehicle ((2-hydroxypropyl)- $\beta$ -cyclodextrin) or one of the two sEHIs: 1-adamantan-3-(5-(2-(2-ethylethoxy)ethoxy)pentyl)urea (AEP; 0.1 mg/ml) or 12-(3-adamantan-1-yl-ureido)-dodecanoic acid (AUDA; 0.1 mg/ml) (Fig. 1A; and see Tables 1–3 and Fig. 7, which are published as supporting information on the PNAS web site; ref. 21) in drinking water after surgery for a period of 3 weeks. The synthesis of both sEHIs was scaled up from described procedures (21). AUDA has relatively low solubility in water compared with AEP and requires (2-hydroxypropyl)- $\beta$ -cyclodextrin to achieve the concentration required for administration in drinking water. Measurement of plasma drug levels is presented in *Supporting Text*, which is published as supporting information on the PNAS web site. Fig. 1B shows photomicrographs of examples of whole hearts from TAC mice treated with AEP or AUDA in the

Author contributions: D.X., N.L., B.D.H., and N.C. designed research; D.X., N.L., Y.H., V.T., L.L., D.T., R.K.P.M., A.S., B.B.D., and N.C. performed research; H.-J.T., I.-H.K., R.L., and B.D.H. contributed new reagents/analytic tools; D.X., N.L., Y.H., V.T., L.L., B.D.H., and N.C. analyzed data; and B.D.H. and N.C. wrote the paper.


Conflict of interest statement: N.C. and B.D.H. have filed patents for the University of California for sEH and cardiac hypertrophy therapy. B.D.H. founded Arête Therapeutics to move sEH inhibitors into clinical trials.

Abbreviations: Ang II, angiotensin II; ANF, atrial natriuretic factor; AEP, 1-adamantan-3-(5-(2-(2-ethylethoxy)ethoxy)pentyl)urea; AUDA, 12-(3-adamantan-1-yl-ureido)-dodecanoic acid; DHET, dihydroxyeicosatrienoic acid; EET, epoxyeicosatrienoic acid; LV-ESD, left ventricular end systolic dimension; MHC, myosin heavy chain; sEH, soluble epoxide hydrolase; sEH, sEH inhibitor; TAC, thoracic aortic constriction.

<sup>¶</sup>To whom correspondence may be addressed. E-mail: bdhammock@ucdavis.edu.

<sup>\*\*</sup>To whom correspondence may be addressed at: Division of Cardiovascular Medicine, University of California, One Shields Avenue, GBSF 6315, Davis, CA 95616. E-mail: nchiamvimonvat@ucdavis.edu.


© 2006 by The National Academy of Sciences of the USA



**Fig. 1.** Inhibition of cardiac hypertrophy in TAC mice by sEHs (AEPU and AUDA). (A) Structure of the two sEHs used in our studies: AEPU or AUDA. (B) Examples of whole hearts from TAC mice treated with AEPU or AUDA compared with TAC alone or sham-operated hearts. The mice were killed after 3 weeks of followup. (Scale, 1 cm.) (C) Analysis of heart/body weight ratios (mg/g) demonstrated that AEPU- and AUDA-free acid inhibited the increase in heart size in TAC mice compared with untreated TAC mice. Error bars are  $\pm$ SEM of  $n = 16$ . (D) Histologic sections (H&E staining) of a sham-operated (control) and TAC mouse hearts, showing cardiac hypertrophy at 3 weeks in the TAC mouse. Treatment of TAC mice with AEPU (0.1 mg/ml) in drinking water prevented the development of cardiac hypertrophy (\*,  $P < 0.05$ ). All histologic sections are presented with the atria on top and the right ventricle to the left. (Scale bars, 200  $\mu$ m.)

drinking water for 3 weeks compared with TAC alone or sham-operated hearts after 3 weeks of followup. All TAC mice showed the expected increase in heart size, with dilatation of all chambers as previously described in this model (23, 24). In contrast, treatment with sEHs prevented development of cardiac hypertrophy. Summary data are shown in Fig. 1C illustrating a significant increase in the ratio of heart weight/body weight (HW/BW) in the TAC group but no significant changes in the HW/BW ratio in the TAC groups treated with AEPU or AUDA compared with sham-treated group were found. The H&E histologic sections in Fig. 1D show sections of a sham-operated and TAC hearts with or without the treatment. The TAC heart showed evidence of cardiac hypertrophy and chamber dilatation. The hypertrophic response was prevented by using either AEPU or AUDA in the drinking water. There were no changes in the sham-operated mice treated with AEPU. Both AEPU and AUDA are potent inhibitors of sEH; however, AUDA is far more lipophilic than AEPU and can be viewed as a possible mimic of 14,15 EET. The fact that these two inhibitors of sEH with radically different physical properties both yield similar biology suggests that the inhibition of sEH is involved.

**Noninvasive Assessment of Cardiac Hypertrophy by Echocardiography.** We assessed the chamber size and wall thickness in sham-operated and TAC mice compared with TAC mice treated with sEHs by using echocardiography. As shown in Fig. 2A using M model echocardiography, AEPU prevented the development of cardiac hypertrophy in TAC mice after 3 weeks of followup. Fig. 2B and C summarize the percentage of fractional shortening and left ventricular end systolic dimension (LV-ESD) from sham, TAC, and TAC mice treated with AEPU or AUDA. Treatment with sEHs prevented the deterioration in fractional shortening and LV-ESD compared with TAC alone. We further analyzed the single isolated cardiac myocytes from the left ventricular free wall from the different groups of mice by using light microscopy. Fig. 2D shows photomicrographs of single ventricular myocytes isolated from the



**Fig. 2.** Assessment of cardiac hypertrophy. (A) Examples of M-mode echocardiography in sham-operated, TAC, and TAC treated with AEPU mice after 3 weeks of treatment, showing evidence of cardiac failure with chamber dilatation in TAC mice. AEPU prevented the development of cardiac hypertrophy and failure in TAC mice. Summary data are shown in B and C. (B) Fractional shortening (FS), a surrogate of systolic function, was calculated from left ventricle dimensions as follows:  $FS = ((EDD - ESD)/EDD) \times 100\%$ . (C) LV-ESD. Data shown are mean  $\pm$  SEM,  $n = 10$ –16 for each group; \*,  $P < 0.05$ . (D) Photomicrographs of single isolated cardiac myocytes from left ventricular free wall. Treatment with compounds AEPU or AUDA prevented hypertrophy of the cardiac myocytes in TAC mice.


four groups of mice. Treatment with AEPU or AUDA prevented the hypertrophy of the cardiac myocytes in TAC mice. Furthermore, the disarray of myofibrils that characterized the hypertrophic hearts was also prevented by treatment with sEHs (Fig. 3A).

**Analysis of Hypertrophic Markers.** To further test the hypothesis that sEHs antagonize initiation of the hypertrophic response, we examined the induction of hypertrophic markers such as atrial natriuretic factor (ANF), skeletal  $\alpha$ -actin, and  $\alpha$  vs.  $\beta$ -myosin heavy chain (MHC) isoforms switch using total RNA isolated from hearts of sham-operated, TAC alone and TAC treated with the sEH (AEPU). Expression of transcripts for ANF, skeletal  $\alpha$ -actin,  $\alpha$ - and  $\beta$ -MHC, and GAPDH were determined. The basal level of these mRNAs was detected in myocytes. Aortic banding resulted in the up-regulation of these hypertrophic markers as described (25). AEPU decreased the transcript expression of the hypertrophic genes including ANF, skeletal  $\alpha$ -actin, and  $\beta$ -MHC as compared with TAC alone (Fig. 3B). These data further support our findings in Figs. 1 and 2 that sEHs have beneficial effects in this *in vivo* model of cardiac hypertrophy. Additional data from microarray analysis are presented in Table 4, which is published as supporting information on the PNAS web site.

**Expression of sEH in Murine Atrial and Ventricular Myocytes.** sEH has been documented to be highly expressed in liver and kidney. To directly document the expression of the enzyme in the heart, we performed Western blot analysis using atrial and ventricular tissues from male mice (C57BL/6J) and polyclonal anti-sEH Ab (1:200 dilution) against human sEH that was raised from rabbits as described (26). As shown in Fig. 3C, both mouse atrial and ventricular tissues expressed a readily detectable level of sEH








**Fig. 4.** Reversal of preestablished cardiac hypertrophy in TAC mice by sEHs. (A) Examples of whole hearts from TAC mice at 6 weeks (TAC 6 week) compared with TAC mice started on the treatment with AEPU at 3 weeks and continued for 3 weeks (TAC 3W+AEPU 3W) or sham-operated hearts (with and without treatment). The mice were killed after 6 weeks of followup. (Scale, 1 cm.) (B) Photomicrograph of the corresponding histologic section (H&E). (Scale bars, 200  $\mu$ m.) (C and D) Summary data for the fractional shortening (C) and LV-ESD (D) from TAC mice at 6 weeks (TAC 6w) compared with TAC mice started on the treatment with AEPU at 3 weeks and continued for 3 weeks (TAC 3w+AEPU 3w) or sham-operated hearts (with and without treatment). The mice were killed after 6 weeks of followup. AEPU significantly reversed the established cardiac hypertrophy in the TAC mice ( $n = 8-10$  for each group) (\*,  $P < 0.05$ ).


mAb (1:200 dilution; Fig. 6A). Treatment with Ang II or PE resulted in nuclear translocation of NF- $\kappa$ B from the cytosol into the nucleus. The effects were inhibited by treatment of the cells with AEPU.

**NF- $\kappa$ B Activation Is Inhibited in the Heart *in Vivo* by sEHs.** The inhibitory effect of sEHs on the activation of NF- $\kappa$ B was further studied in the TAC model (Fig. 6B). Previous studies documented the activation of NF- $\kappa$ B in the murine model of cardiac hypertrophy with aortic banding (6, 8, 12). Here, we tested the hypothesis that sEHs can block the activation of NF- $\kappa$ B in the *in vivo* model using Western blot analysis (Fig. 6B). Total I $\kappa$ B $\alpha$  and phospho-I $\kappa$ B $\alpha$  ( $p$ -I $\kappa$ B $\alpha$ ) levels after TAC were assessed by using a total homogenate from the left ventricular free wall obtained from TAC mice compared with TAC mice treated with AEPU at different time points (0, 1/2, 1, 3, and 6 h after aortic banding). Mice received



**Fig. 6.** Prevention of nuclear translocation of NF- $\kappa$ B by treatment with AEPU in primary mouse neonatal cardiomyocytes. (A) Cardiomyocytes grown on coverslips were serum-starved for 24 h before treatment with Ang II (50  $\mu$ M) or phenylephrine (PE; 50  $\mu$ M) for 45 min or left untreated (control) or Ang II + AEPU (20  $\mu$ M) or PE + AEPU (20  $\mu$ M). Cells were fixed, blocked, and stained with anti-p65 mAb, followed by anti-mouse Texas red-conjugated secondary Ab (red). The nuclear translocation of p65 was visualized by DAPI stain (blue) using a Zeiss confocal microscope. (B) AEPU inhibits NF- $\kappa$ B signaling in the hearts. Total I $\kappa$ B $\alpha$  and phospho-I $\kappa$ B $\alpha$  ( $p$ -I $\kappa$ B $\alpha$ ) levels after ascending aortic constriction (TAC). Proteins (50  $\mu$ g per lane) from left ventricular free wall from banded mice compared with banded mice treated with AEPU at different time points (0, 1/2, 1, 3, and 6 h) were separated by SDS/PAGE. Total I $\kappa$ B $\alpha$  (top blot) and phospho-I $\kappa$ B $\alpha$  (second blot) were detected by immunoblotting. P65 level in the nuclear fraction is shown in the third blot. GAPDH was used as internal loading control. AEPU prevented the increase in the  $p$ -I $\kappa$ B $\alpha$  level and the nuclear translocation of NF- $\kappa$ B.

AEPU (10 mg/kg) vs. (2-hydroxypropyl)- $\beta$ -cyclodextrin (vehicle only; 5 mg/kg) i.p. before surgery. Anti-I $\kappa$ B $\alpha$  and phospho-I $\kappa$ B $\alpha$  mAbs were used (1:200 dilution). The level of NF- $\kappa$ B was further assessed in the nuclear fraction by using anti-p65 Ab (1:200 dilution). There was a progressive increase in the phospho-I $\kappa$ B $\alpha$  and a decrease in the total I $\kappa$ B $\alpha$  level, associated with nuclear translocation of NF- $\kappa$ B, which peaked at  $\approx 3$  h after aortic banding similar to a study documenting the activation of NF- $\kappa$ B in this *in vivo* model (8). Moreover, as shown in Fig. 6B, inhibition of sEH by AEPU prevented the increase in the  $p$ -I $\kappa$ B $\alpha$  level as well as the nuclear translocation of NF- $\kappa$ B induced by aortic banding.



**Fig. 5.** *In vitro* assessment of the effects of sEHs on the development of cardiac hypertrophy. (A) Photomicrographs of primary neonatal mouse cardiac myocytes treated with Ang II for 36 h to induce cardiac hypertrophy or AngII in the presence of AEPU or EET for 36 h. Cells were then fixed and treated with anti- $\alpha$ -actinin2 Abs and secondary Ab conjugated to Texas red. Shown are control neonatal cardiac myocytes and cells treated with Ang II alone (Upper Left and Upper Right, respectively). (Lower Left and Lower Right) Cells treated with Ang II plus AEPU or Ang II plus EET, respectively. (B) Quantification of cell surface area comparing control, AngII, and AngII treated with AEPU at 48 h (50  $\mu$ M). (\*,  $P < 0.05$ ). (C) Dose-response curve of cell surface area in various concentrations of AEPU.



PNAS | December 5, 2006 | vol. 103 | no. 49 | 18737

hypertrophy in this model because a persistent chronic increase in pressure is exerted on the heart in this model by the constriction of the thoracic aorta. Thus, future work is required to directly compare the effects of sEHs and blood pressure-lowering agents that act through separate mechanisms on cardiac hypertrophy in this model.

Finally, we were able to show that use of sEHs can prevent the development of ventricular arrhythmias that occur in association with cardiac hypertrophy. Indeed, cardiac hypertrophy and failure are associated with a significant increase in the risk of arrhythmias and sudden cardiac death (46). Moreover, treatment with conventional antiarrhythmic drugs generally carries a high risk of proarrhythmia and results in a decrease in patient survival (47). The most accepted form of therapy involves implantation of defibrillators, which is invasive and costly (48). Our study has important clinical significance in providing evidence for an alternative strategy in the treatment of cardiac hypertrophy and the prevention of ventricular arrhythmias associated with cardiac hypertrophy.

## Materials and Methods

For more detail, see *Supporting Materials and Methods*, which is published as supporting information on the PNAS web site.

**Compensated Cardiac Hypertrophy Model in Mice.** TACs were performed in 8-week-old male C57BL/6J mice as described (24). Animals with TAC were randomized to treatment with two different sEHs: AEPu in drinking water (0.1 mg/ml), AUDA-free acid (0.1 mg/ml) in drinking water, or vehicle only for 3 weeks. Sham-operated animals received the same surgery except that the ligature was not tied. Echocardiograms to assess cardiac function were performed at the end of the experiments by using M-mode and 2D measurements. *In vivo* EPS were performed as described (31, 49).

Standard pacing protocols were used. Each animal underwent an identical pacing and programmed stimulation protocol.

**Semiquantitative RT-PCR Analysis.** Total RNA from left ventricular free wall was reverse-transcribed. cDNA was then diluted with water to generate serial dilutions (1:1, 1:2, and 1:10) and used for PCR amplification with gene-specific primers.

**Western Blot Analysis.** Immunoblots were performed as described (50). The following primary Abs were used: (i) anti- $\text{I}\kappa\text{B}\alpha$  and phosphor- $\text{I}\kappa\text{B}\alpha$  mAbs (1:200 dilution, Santa Cruz Biotechnology, Santa Cruz, CA); (ii) anti-p65 (Santa Cruz Biotechnology, 1:200 dilution); (iii) polyclonal anti-sEH Ab (1:2,000) against human sEH was raised from rabbits as described (26); and (iv) anti-GAPDH Ab (Sigma) was used as an internal loading control.

**Immunofluorescence Confocal Microscopy.** Immunofluorescence labeling was performed as described and imaged using a confocal laser scanning microscopy (50) (Pascal Zeiss, Thornwood, NY). Control experiments performed with secondary Ab only did not show positive staining under the same experimental conditions used. Identical settings were used for all of the specimens.

We acknowledge the University of California, Davis, Health System (UCDHS) Confocal Microscopy Facility. This work was supported by a Department of Veterans Affairs Merit Review Grant and National Institutes of Health Grants HL68507 and HL67737 (to N.C.) and ES11634 (to B.B.D.). Partial support was provided by National Institute on Environmental Health Sciences (NIEHS) Grant R37 ES02710, NIEHS Superfund Basic Research Program Grant P42 ES04699, NIEHS Center Grant P30 ES05707, a Technology Translational Grant from UCDHS (to B.D.H.), and the Howard Hughes Medical Institute Med into Grad Initiative of the Graduate Science Education Program (to University of California, Davis).

1. Frey N, Olson EN (2003) *Annu Rev Physiol* 65:45–79.
2. Molkenkin J, Dorn GW, II (2001) *Annu Rev Physiol* 63:391–426.
3. Frey N, Katus HA, Olson EN, Hill JA (2004) *Circulation* 109:1580–1589.
4. Morisco C, Sadoshima J, Trimarco B, Arora R, Vatner DE, Vatner SF (2003) *Am J Physiol Heart Circ Physiol* 284:H1043–H1047.
5. Purcell NH, Tang G, Yu C, Mercurio F, DiDonato JA, Lin A (2001) *Proc Natl Acad Sci USA* 98:6668–6673.
6. Hirotani S, Otsu K, Nishida K, Higuchi Y, Morita T, Nakayama H, Yamaguchi O, Mano T, Matsumura Y, Ueno H, et al. (2002) *Circulation* 105:509–515.
7. Gupta S, Purcell NH, Lin A, Sen S (2002) *J Cell Biol* 159:1019–1028.
8. Cook SA, Novikov MS, Ahn Y, Matsui T, Rosenzweig A (2003) *Circulation* 108:664–667.
9. Purcell NH, Molkenkin JD (2003) *Circulation* 108:638–640.
10. Freund C, Schmidt-Ullrich R, Baurand A, Dunger S, Schneider W, Loser P, El-Jamali A, Dietz R, Scheidereit C, Bergmann MW (2005) *Circulation* 111:2319–2325.
11. Kawano S, Kubota T, Monden Y, Kawamura N, Tsutsui H, Takeshita A, Sunagawa K (2005) *Cardiovasc Res* 67:689–698.
12. Li Y, Ha T, Gao X, Kelley J, Williams D L, Browder IW, Kao R L, Li C (2004) *Am J Physiol Heart Circ Physiol* 287:H1712–H1720.
13. Karin M, Lin A (2002) *Nat Immunol* 3:221–227.
14. van Empel VP, De Windt L J (2004) *Cardiovasc Res* 63:487–499.
15. Yu Z, Xu F, Huse LM, Morisseau C, Draper AJ, Newman JW, Parker C, Graham L, Engler MM, Hammock BD, et al. (2000) *Circ Res* 87:992–998.
16. Zeldin DC, Moomaw CR, Jesse N, Tomer KB, Beetham J, Hammock BD, Wu S (1996) *Arch Biochem Biophys* 330:87–96.
17. Spector AA, Fang X, Snyder GD, Weintraub NL (2004) *Prog Lipid Res* 43:55–90.
18. Campbell WB (2000) *Trends Pharmacol Sci* 21:125–127.
19. Node K, Huo Y, Ruan X, Yang B, Spiecker M, Ley K, Zeldin DC, Liao JK (1999) *Science* 285:1276–1279.
20. Imig JD, Zhao X, Capdevila JH, Morisseau C, Hammock BD (2002) *Hypertension* 39:690–694.
21. Morisseau C, Hammock BD (2005) *Annu Rev Pharmacol Toxicol* 45:311–333.
22. Schmelzer KR, Kubala L, Newman JW, Kim IH, Eiserich JP, Hammock BD (2005) *Proc Natl Acad Sci USA* 102:9772–9777.
23. Takimoto E, Champion HC, Li M, Belardi D, Ren S, Rodriguez ER, Bedja D, Gabrielson KL, Wang Y, Kass DA (2005) *Nat Med* 11:214–222.
24. Tarnavski O, McMullen JR, Schinke M, Nie Q, Kong S, Izumo S (2004) *Physiol Genomics* 16:349–360.
25. Chien KR, Knowlton KU, Zhu H, Chien S (1991) *FASEB J* 5:3037–3046.
26. Wixtrom RN, Silva MH, Hammock BD (1988) *Placenta* 9:559–563.
27. Li PL, Campbell WB (1997) *Circ Res* 80:877–884.
28. Karin M (1999) *J Biol Chem* 274:27339–27342.
29. Karin M (1999) *Oncogene* 18:6867–6874.
30. Rothwarf DM, Karin M (1999) *Sci STKE* 1999:RE1.
31. Zhang Z, He Y, Tuteja D, Xu D, Timofeyev V, Zhang Q, Glatter KA, Xu Y, Shin HS, Low R, Chiamvimonvat N (2005) *Circulation* 112:1936–1944.
32. Li H, Kedar V, Zhang C, McDonough H, Arya R, Wang D, Z., Patterson C. (2004) *J Clin Invest* 114:1058–1071.
33. Antos CL, McKinsey TA, Dreitz M, Hollingsworth LM, Zhang CL, Schreiber K, Rindt H, Gorczynski RJ, Olson EN (2003) *J Biol Chem* 278:28930–28937.
34. Czubrynt MP, Olson EN (2004) *Recent Prog Horm Res* 59:105–124.
35. Molkenkin JD, Olson EN (1997) *Circulation* 96:3833–3835.
36. Rothermel BA, McKinsey TA, Vega RB, Nicol RL, Mammen P, Yang J, Antos CL, Shelton JM, Bassel-Duby R, Olson EN, Williams RS (2001) *Proc Natl Acad Sci USA* 98:3328–3333.
37. Sussman MA, Lim HW, Gude N, Taigen T, Olson EN, Robbins J, Colbert MC, Gualberto A, Wiczorek DF, Molkenkin JD (1998) *Science* 281:1690–1693.
38. Baeuerle PA, Baltimore D (1996) *Cell* 87:13–20.
39. Matsumori A, Sasayama S (2001) *Heart Fail Rev* 6:129–136.
40. Wong SC, Fukuchi M, Melnyk P, Rodger I, Giaid A (1998) *Circulation* 98:100–103.
41. Flemming I (2001) *Circ Res* 89:753–762.
42. Roman RJ, Maier KG, Sun CW, Harder DR, Alonso-Galicia M (2000) *Clin Exp Pharmacol Physiol* 27:855–865.
43. Zeldin DC, Wei S, Falck JR, Hammock BD, Snapper JR, Capdevila JH (1995) *Arch Biochem Biophys* 316:443–451.
44. Esposito G, Rapacciuolo A, Naga Prasad SV, Takaoka H, Thomas SA, Koch WJ, Rockman HA (2002) *Circulation* 105:85–92.
45. Sano M, Schneider MD (2002) *Circulation* 105:8–10.
46. Hart G (1993) *J R Coll Physicians London* 27:429–434.
47. Ruskin JN (1989) *N Engl J Med* 321:386–388.
48. Moss AJ, Zareba W, Hall WJ, Klein H, Wilber DJ, Cannom DS, Daubert JP, Higgins SL, Brown MW, Andrews ML (2002) *N Engl J Med* 346:877–883.
49. Berul CI, Aronovitz MJ, Wang PJ, Mendelsohn ME (1996) *Circulation* 94:2641–2648.
50. Xu Y, Zhang Z, Timofeyev V, Sharma D, Xu D, Tuteja D, Dong PH, Ahmed GU, Ji Y, Shull GE. (2005) *J Physiol* 562:745–758.